

JAMIE SANDS, D.D.S.

COSMETIC, RESTORATIVE AND PREVENTIVE CARE

ORAL ANTI-AGING™

Patient Post-Op Instructions - Scaling and Root Planing

To minimize the discomfort and encourage proper healing following your scaling and root planing, follow these instructions:

- After the procedure, take aspirin, acetaminophen (Tylenol®), or ibuprofen (Advil®) before the anesthetic wears off. Continue to take up to 3 tabs or 600 mgs Ibuprofen and 2 extra strength Tylenol every four hours for the next two days.
- A saltwater solution (1/2 teaspoon salt + 1/2 teaspoon soda + 8 ounces warm water) swished in your mouth for 2 to 3 minutes every hour may make your mouth more comfortable.
- Use a soft toothbrush at least two times a day. Be gentle and clean thoroughly. Slight bleeding may occur while brushing as the tissues begin to heal.
- Avoid strong spicy seasonings and hard crunchy foods for the next few days.
- Smoking should be stopped. Success of the treatment will be substantially reduced by cigarette smoke chemicals in your body.
- As the tissues heal, some temporary sensitivity to cold may occur. Use a desensitizing toothpaste (such as Sensodyne®), or fluoride gel (such as Prevident® or Gel-Kam®) frequently (at least 4 times/day) for 1 to 2 weeks. Also, the cleaner the teeth are kept, the less sensitive they will be.
- Faithfully use any other oral hygiene aids that have been recommended (floss, Perio-Aid®, rubber tip, Sonicare®, Proxabrush®, Gel-Kam® fluoride, Peridex® mouthrinse, etc).
- If you have any questions or concerns, please call our office at **818-766-6767**